Barrettine Divergendent Health Divergendent He

SPECIES SPECIFIC AGAINST: PEST BIRDS

Preparing surfaces for treatment

Ensure appropriate PPE is used when preparing the site, removal of all nests and other debris. For effective installation of **Ornaway Bird Gel** all surfaces within the installation area must be thoroughly cleaned and free from faecal and organic matter.

Enhanced efficacy will be gained by treating the cleaned areas with a fine mist spray of a proprietary biocide (eg, Guanaway). Do not over-wet as this will lengthen drying time and inhibit adhesion of silicone. **Ornaway Bird Gel** will only be completely effective if the site is properly cleaned before installation.

APPLICATION

Cut off cartridge seal, cut nozzle to the required size and apply 15 g - 17 g of **Ornaway Bird Gel** into each dish, **Ornaway Bird Gel** can also be applied direct to some surfaces (gel may stain some surfaces).

STORAGE

Cool dry conditions. Keep out of reach of children.

DISPOSAL

Product contains food and pharmaceutical grade materials and can be disposed of appropriately on site or at an approved waste disposal facility. Do not reuse the caulking tube, once empty please recycle cartridge (if available) or dispose of with normal / household waste. Never dispose unused product down any indoor or outdoor drains.



Developed by UK scientists. Manufactured in the UK.

Ornaway Bird Gel does not contain polybutene.

Barrettine Environmental Health St. Ivel Way, Warmley, Bristol BS30 8TY. T: +44 (0) 117 967 2222 F: +44 (0) 117 961 4122 E: beh@barrettine.co.uk www.barrettine.co.uk

SPACING & APPLICATION RATES

High pressure site 4-6 inches (10-15cm)

Medium pressure site 8-10 inches (20-25cm)

Low pressure site 2-20 inches (30-50cm)

